Complete Summary

GUIDELINE TITLE

Stress-related conditions.

BIBLIOGRAPHIC SOURCE(S)

Stress-related conditions. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2004. 27 p. [129 references]

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Harris, J, ed. Occupational Medicine Practice Guidelines: American College of Occupational and Environmental Medicine. Beverly Farms, MA: OEM Press; 1997.

** REGULATORY ALERT **

FDA WARNING/REGULATORY ALERT

Note from the National Guideline Clearinghouse: This guideline references a drug(s) for which important revised regulatory information has been released.

May 2, 2007, Antidepressant drugs: Update to the existing black box warning
on the prescribing information on all antidepressant medications to include
warnings about the increased risks of suicidal thinking and behavior in young
adults ages 18 to 24 years old during the first one to two months of
treatment.

COMPLETE SUMMARY CONTENT

** REGULATORY ALERT **

SCOPE

METHODOLOGY - including Rating Scheme and Cost Analysis

RECOMMENDATIONS

EVIDENCE SUPPORTING THE RECOMMENDATIONS

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS QUALIFYING STATEMENTS

IMPLEMENTATION OF THE GUIDELINE

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IDENTIFYING INFORMATION AND AVAILABILITY DISCLAIMER

DISEASE/CONDITION(S)

Stress-related conditions

GUIDELINE CATEGORY

Diagnosis Evaluation Management Treatment

CLINICAL SPECIALTY

Family Practice Internal Medicine Physical Medicine and Rehabilitation Preventive Medicine Psychiatry

INTENDED USERS

Advanced Practice Nurses Physician Assistants Physicians Utilization Management

GUIDELINE OBJECTIVE(S)

- To provide information and guidance on generally accepted elements of quality care in occupational and environmental medicine
- To improve the efficiency with which the diagnostic process is conducted, the specificity of each diagnostic test performed, and the effectiveness of each treatment in relieving symptoms and achieving cure
- To help occupational physicians and primary care practitioners manage employed patients with acute stress-related conditions of relatively short duration
- To help clinicians identify patients who need urgent referral for psychiatric care and provide a framework for treating the majority of patients who do not

TARGET POPULATION

Adults with potentially work-related stress-related conditions seen in primary care settings

INTERVENTIONS AND PRACTICES CONSIDERED

Evaluation

- 1. Medical history including symptoms, stressors, coping mechanisms, and other resources
- 2. Physical examination including a mental status examination

Diagnosis

- 1. Evaluating for potentially life-threatening or other serious diseases that the history and physical examination may suggest
- 2. Coding the visit appropriately

Treatment/Management

- 1. Patient education
- 2. Specialty referral
- 3. Management of medical conditions
- 4. Modification of maladaptive coping mechanisms
- 5. Aerobic exercise
- 6. Stress management techniques
 - Relaxation techniques
 - Behavioral techniques
 - Cognitive techniques and therapy
 - Stress inoculation therapy
- 7. External resources and referrals
 - Employee assistance programs
- 8. Pharmacotherapy (limited use or in conjunction with specialty referral)
 - Anxiolytics
 - Antidepressants
 - Antipsychotics
- 9. Modified work and accommodations
- 10. Organizational interventions
- 11. Determining disability duration
- 12. Follow-up, as appropriate

MAJOR OUTCOMES CONSIDERED

Missed work days

METHODOLOGY

METHODS USED TO COLLECT/SELECT EVIDENCE

Searches of Electronic Databases

DESCRIPTION OF METHODS USED TO COLLECT/SELECT THE EVIDENCE

Note from the National Guideline Clearinghouse (NGC): The American College of Occupational and Environmental Medicine contracted the Work Loss Data Institute to provide medical library research services.

NUMBER OF SOURCE DOCUMENTS

Not stated

METHODS USED TO ASSESS THE QUALITY AND STRENGTH OF THE EVIDENCE

Expert Consensus

RATING SCHEME FOR THE STRENGTH OF THE EVIDENCE

Not applicable

METHODS USED TO ANALYZE THE EVIDENCE

Review

Review of Published Meta-Analyses

DESCRIPTION OF THE METHODS USED TO ANALYZE THE EVIDENCE

Contributors reviewed at least one chapter each and reviewed the relevant medical literature that had been published since the creation of the original Guidelines in 1997.

METHODS USED TO FORMULATE THE RECOMMENDATIONS

Expert Consensus

DESCRIPTION OF METHODS USED TO FORMULATE THE RECOMMENDATIONS

Not stated

RATING SCHEME FOR THE STRENGTH OF THE RECOMMENDATIONS

Not applicable

COST ANALYSIS

A formal cost analysis was not performed and published cost analyses were not reviewed.

METHOD OF GUIDELINE VALIDATION

Internal Peer Review

DESCRIPTION OF METHOD OF GUIDELINE VALIDATION

Following the chapter and literature review, participants provided written or verbal comments to the American College of Occupational and Environmental Medicine's Practice Guidelines Committee.

Verbal comments were in the form of participation in multi-specialty conference calls, during which the issues raised in each chapter were extensively discussed. Draft chapters were prepared and distributed by the American College of Occupational and Environmental Medicine to all chapter reviewers. Follow-up multi-specialty teleconferences were then held as appropriate, during which time the draft was again reviewed.

RECOMMENDATIONS

MAJOR RECOMMENDATIONS

General Approach and Basic Principles (refer to the original guideline document for more detailed information)

Stressors may be any life event or circumstance that exerts a physical, emotional, or cognitive demand on the individual. The lifetime prevalence of major stressful life events is 100%, so associated stress-related symptoms may be considered a normal condition of human existence.

- Stress is not a diagnosis, disease, or syndrome. It is a nonspecific set of
 emotions or physical symptoms that may or may not be associated with a
 disease or syndrome. Whether or not stress contributes to a disease or
 syndrome depends on the vulnerability of the individual; the intensity,
 duration, and meaning of the stress; and the nature and availability of
 modifying resources.
- The initial assessment of patients with acute stress-related conditions focuses on detecting potentially serious psychopathology, or red flag conditions, requiring urgent specialty referral. The majority of patients with stress-related conditions will not have red flags and can be safely and effectively managed by occupational or primary care physicians.
- Relief of stress depends on its precipitants, which are often multifactorial.
 Psychosocial, workplace, or socioeconomic issues can be explored with the patient to facilitate early identification of precipitating factors and appropriate interventions that may prevent delayed recovery or relapse. An open, honest discussion of the underlying factors often results in an increase in the patient's insight and coping skills, which itself helps alleviate many stress-related symptoms.
- Worksite interventions may be helpful in mitigating or eliminating inciting stressors, depending on the source of the complaints.
- Medications generally have a limited role. Limit use of anti-anxiety agents to short periods of time (i.e., periods when overwhelming anxiety limits the patient's ability to work or effectively perform the activities of daily living). Antidepressant or antipsychotic medication may be prescribed for major depression or psychosis; however, this is best done in conjunction with specialty referral.
- Patients are encouraged to enhance their individual coping skills and to decrease or discontinue maladaptive coping mechanisms such as excessive use of alcohol, tobacco, or other drugs, or excessive food intake. Patients are counseled to redirect their energy to regular aerobic exercise, relaxation techniques, and cognitive coping mechanisms.

- For uncomplicated cases, absence from work should not exceed one work week. Referral for mental health professional assessment may be considered for patients whose anticipated absence from work will exceed one week.
- If symptoms become disabling despite primary care interventions or persist beyond three months, referral to a mental health professional is indicated.

CLINICAL ALGORITHM(S)

None provided

EVIDENCE SUPPORTING THE RECOMMENDATIONS

TYPE OF EVIDENCE SUPPORTING THE RECOMMENDATIONS

The type of supporting evidence is not specifically stated for each recommendation.

BENEFITS/HARMS OF IMPLEMENTING THE GUIDELINE RECOMMENDATIONS

POTENTIAL BENEFITS

- Improved efficiency of the diagnostic process
- Effective treatment resulting in symptom alleviation and cure

POTENTIAL HARMS

- Anxiolytics can lead to dependence and do not alter stressors or the individual's coping mechanisms.
- Antidepressants have many side effects and can result in decreased work performance or mania in some people.
- Antipsychotics can decrease motivation and effectiveness at work.

QUALIFYING STATEMENTS

QUALIFYING STATEMENTS

The American College of Occupational and Environmental Medicine provides this segment of guidelines for practitioners and notes that decisions to adopt particular courses of actions must be made by trained practitioners on the basis of the available resources and the particular circumstances presented by the individual patient. Accordingly, the American College of Occupational and Environmental Medicine disclaims responsibility for any injury or damage resulting from actions taken by practitioners after considering these guidelines.

IMPLEMENTATION OF THE GUIDELINE

DESCRIPTION OF IMPLEMENTATION STRATEGY

An implementation strategy was not provided.

INSTITUTE OF MEDICINE (IOM) NATIONAL HEALTHCARE QUALITY REPORT CATEGORIES

IOM CARE NEED

Getting Better

IOM DOMAIN

Effectiveness Patient-centeredness

IDENTIFYING INFORMATION AND AVAILABILITY

BIBLIOGRAPHIC SOURCE(S)

Stress-related conditions. Elk Grove Village (IL): American College of Occupational and Environmental Medicine (ACOEM); 2004. 27 p. [129 references]

ADAPTATION

Not applicable: The guideline was not adapted from another source.

DATE RELEASED

1997 (revised 2004)

GUIDELINE DEVELOPER(S)

American College of Occupational and Environmental Medicine - Medical Specialty Society

SOURCE(S) OF FUNDING

American College of Occupational and Environmental Medicine

GUIDELINE COMMITTEE

American College of Occupational and Environmental Medicine Practice Guidelines Committee

COMPOSITION OF GROUP THAT AUTHORED THE GUIDELINE

Chapter Lead: Lee S. Glass, MD

Committee Members: Jennifer H. Christian, MD, MPH, FACPM; Philip I. Harber, MD, MPH, FACOEM, FCCP; John P. Holland, MD, MPH, FACOEM; Kathryn L. Mueller, MD, MPH, FACEP, FACOEM; Douglas J. Patron, MD, MSPH; Bernyce M. Peplowski, DO, MS; and Jack Richman, MD, CCFP, DOHS, FACOEM

Timothy J. Key, MD, MPH, FACOEM, as Responsible Officer and ACOEM President Elect, and Edward A. Emmett, MD, MS, FACOEM, Chair of the ACOEM Council on Occupational and Environmental Medical Practice, contributed to the development of the guidelines as well.

FINANCIAL DISCLOSURES/CONFLICTS OF INTEREST

Not stated

GUIDELINE STATUS

This is the current release of the guideline.

This guideline updates a previous version: Harris, J, ed. Occupational Medicine Practice Guidelines: American College of Occupational and Environmental Medicine. Beverly Farms, MA: OEM Press; 1997.

GUIDELINE AVAILABILITY

Print copies are available from ACOEM, 25 Northwest Point Boulevard, Suite 700, Elk Grove Village, IL 60007; Phone: 847-818-1800 x399. To order a subscription to the online version, call 800-441-9674 or visit http://www.acoempracquides.org/.

AVAILABILITY OF COMPANION DOCUMENTS

None available

PATIENT RESOURCES

None available

NGC STATUS

This NGC summary was completed by ECRI on May 31, 2006. The information was verified by the guideline developer on November 3, 2006. This summary was updated by ECRI Institute on November 9, 2007, following the U.S. Food and Drug Administration advisory on Antidepressant drugs.

COPYRIGHT STATEMENT

The American College of Occupational and Environmental Medicine, the signator of this license, represent and warrant that they are the publisher of the guidelines and/or possess all rights necessary to grant the license rights to AHRQ and its agents.

DISCLAIMER

NGC DISCLAIMER

The National Guideline Clearinghouse[™] (NGC) does not develop, produce, approve, or endorse the guidelines represented on this site.

All guidelines summarized by NGC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public or private organizations, other government agencies, health care organizations or plans, and similar entities.

Guidelines represented on the NGC Web site are submitted by guideline developers, and are screened solely to determine that they meet the NGC Inclusion Criteria which may be found at http://www.quideline.gov/about/inclusion.aspx.

NGC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or clinical efficacy or effectiveness of the clinical practice guidelines and related materials represented on this site. Moreover, the views and opinions of developers or authors of guidelines represented on this site do not necessarily state or reflect those of NGC, AHRQ, or its contractor ECRI Institute, and inclusion or hosting of guidelines in NGC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding guideline content are directed to contact the guideline developer.

© 1998-2008 National Guideline Clearinghouse

Date Modified: 10/6/2008

